(I)

Although certain embodiments are described in detail above, those having ordinary skill in the art will clearly understand that many modifications are possible in the claims without departing from the teachings thereof. All such modifications are intended to be encompassed within 5 the claims of the invention.

Embodiments of the Invention

A1. A pharmaceutical composition comprising an effective amount of a compound of the formula:

$$R^{7}$$
 $N$ 
 $N$ 
 $N$ 
 $N$ 
 $R^{4}$ 
 $O$ 
 $O$ 
 $O$ 
 $R^{2}$ 
 $O$ 
 $O$ 
 $R^{3}$ 

wherein R1 and R2 are independently selected from H,  $\rm C_1\text{-}C_6$  alkyl,  $\rm C_1\text{-}C_6$  substituted alkyl,  $\rm C_6\text{-}C_{20}$  aryl,  $\rm C_6\text{-}C_{20}$ substituted aryl,  $C_6$ - $C_{20}$  arylalkyl,  $C_6$ - $C_{20}$  substituted arylalkyl, acyloxymethyl esters  $-CH_2OC(=O)R^9$  and acyloxymethyl carbonates —CH<sub>2</sub>OC(=O)OR<sup>9</sup> where R<sup>9</sup> is <sup>30</sup>  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  substituted alkyl,  $C_6$ - $C_{20}$  aryl and  $C_6$ - $C_{20}$ 

 $\rm R^3$  is selected from H,  $\rm C_1\text{-}C_6$  alkyl,  $\rm C_1\text{-}C_6$  substituted alkyl, or  $\rm CH_2OR^8$  where  $\rm R^8$  is  $\rm C_1\text{-}C_6$  alkyl,  $\rm C_1\text{-}C_6$  hydroxyalkyl and  $C_1$ - $C_6$  haloalkyl;  $R^4$  and  $R^5$  are independently selected from H, NH<sub>2</sub>, NHR

and NR<sub>2</sub> where R is C<sub>1</sub>-C<sub>6</sub> alkyl; and

 $R^6$  and  $R^7$  are independently selected from H and  $C_1$ - $C_6$ alkyl;

or a physiologically functional derivative thereof;

in combination with an effective amount of a compound of the formula

$$B$$
 (2) 45

wherein B is selected from adenine, guanine, cytosine, uracil, thymine, 7-deazaadenine, 7-deazaguanine, 7-deaza-8-azaguanine, 7-deaza-8-azaadenine, inosine, nebularine, nitropyrrole, nitroindole, 2-aminopurine, 2-amino-6-chloropurine, 2,6-diaminopurine, hypoxanthine, pseudouridine, 55 J 5-fluorocytosine, 5-chlorocytosine, 5-bromocytosine, 5-iodocytosine, pseudocytosine, pseudoisocytosine, 5-propynylcytosine, isocytosine, isoguanine, 7-deazaguanine, 2-thiopyrimidine, 6-thioguanine, 4-thiothymine, 4-thiouracil, O<sup>6</sup>-methylguanine, N<sup>6</sup>-methyladenine, O<sup>4</sup>-methylthymine, 60 5,6-dihydrothymine, 5,6-dihydrouracil, 4-methylindole, and a pyrazolo[3,4-D]pyrimidine; and

R is selected from H,  $C_1$ - $C_{18}$  alkyl,  $C_1$ - $C_{18}$  substituted alkyl,  $C_2$ - $C_{18}$  alkenyl,  $C_2$ - $C_{18}$  substituted alkenyl,  $C_2$ - $C_{18}$ alkynyl,  $C_2$ - $C_{18}$  substituted alkynyl,  $C_6$ - $C_{20}$  aryl,  $C_6$ - $C_{20}$  65 substituted aryl,  $C_2$ - $C_{20}$  heterocycle,  $C_2$ - $C_{20}$  substituted heterocycle, phosphonate, phosphophosphonate, diphospho-

phosphonate, phosphate, diphosphate, triphosphate, polyethyleneoxy or a physiologically functional derivative thereof; and a pharmaceutically acceptable carrier.

B2. A composition of embodiment A1 wherein, in formula 1, R1 and R2 are independently selected from H, C1-C6 alkyl,  $C_1$ - $C_6$  substituted alkyl,  $C_6$ - $C_{20}$  aryl,  $C_6$ - $C_{20}$  substituted aryl,  $C_6$ - $C_{20}$  arylalkyl,  $C_6$ - $C_{20}$  substituted arylalkyl, acyloxymethyl esters — $CH_2OC(=O)R^9$  and acyloxymethyl carbonates — $CH_2OC(=O)OR^{\circ}$  where  $R^{\circ}$  is C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> substituted alkyl, C<sub>6</sub>-C<sub>20</sub> aryl and  $C_6$ - $C_{20}$  substituted aryl; and  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are independently H or C<sub>1</sub>-C<sub>6</sub> alkyl.

C3. A composition of embodiment A1 wherein, in formula 2, B is cytosine or a 5-halocytosine.

15 D4. A composition of embodiment A1 wherein, in formula 1, R1 and R2 are independently selected from H, C1-C6 alkyl,  $C_1$ - $C_6$  substituted alkyl,  $C_6$ - $C_{20}$  aryl,  $C_6$ - $C_{20}$  substituted aryl,  $C_6$ - $C_{20}$  arylalkyl,  $C_6$ - $C_{20}$  substituted arylalkyl, acyloxymethyl esters — $CH_2OC(=O)R^9$  and acyloxymethyl carbonates —CH<sub>2</sub>OC(=O)OR<sup>9</sup> where R<sup>9</sup> is  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  substituted alkyl,  $C_6$ - $C_{20}$  aryl and  $C_6$ - $C_{20}$  substituted aryl; and  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are independently H or  $C_1$ - $C_6$  alkyl; and, in formula 2, B is cytosine or a 5-halocytosine.

25 E5. A composition of embodiment D4 wherein, in formula 1, R1 and R2 are independently selected from H, acyloxymethyl esters —CH<sub>2</sub>OC(=O)R<sup>9</sup> and acyloxymethyl carbonates — $CH_2OC(=O)OR^9$  where  $R^9$  is  $C_1$ - $C_6$  alkyl; and R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> are independently H or C<sub>1</sub>-C<sub>6</sub> alkyl; and, in formula 2, B is cytosine or a 5-halocytosine and R is H.

F6. A composition of embodiment E5 wherein, in formula 1, R<sup>1</sup> and R<sup>2</sup> are independently selected from H and —CH<sub>2</sub>OC(=O)OCH(CH<sub>3</sub>)<sub>2</sub>; R<sup>3</sup> is —CH<sub>3</sub>; and R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> are H; and, in formula 2, B is 5-fluorocytosine

G7. A pharmaceutical composition comprising a pharmaceutically effective amount of [2-(6-amino-purin-9-yl)-1methyl-ethoxymethyl]-phosphonic acid diisopropoxycarbonyloxymethyl ester fumarate (tenofovir disoproxil fumarate) or a physiologically functional derivative thereof and a pharmaceutically effective amount of (2R, 5S,)-4-amino-5-fluoro-1-(2-hydroxymethyl-1,3-oxathiolan-5-yl)-(1H)-pyrimidin-2-one (emtricitabine) or a physiologically functional derivative thereof; and a pharmaceutically acceptable carrier.

H8. A pharmaceutical formulation of embodiment A1 to G7 further comprising a third active ingredient selected from the group consisting of a protease inhibitor, a nucleoside or nucleotide reverse transcriptase inhibitor, a nonnucleoside reverse transcriptase inhibitor, and an integrase inhibitor.

19. A pharmaceutical formulation of embodiments A1 to H8 in unit dosage form.

10. A method for the treatment or prevention of the symptoms or effects of an HIV infection in an infected animal which comprises administering to said animal a pharmaceutical composition of embodiments claims A1 to I9. We claim:

1. A fixed dose combination pharmaceutical dosage form comprising 300 mg tenofovir disoproxil fumarate and 200 mg emtricitabine; a binder selected from the group consisting of povidone, gelatin, hydroxypropyl methylcellulose, cellulose, microcrystalline cellulose, starch, and acacia; a disintegrant selected from sodium starch glycolate, crosslinked-povidone, cross-linked sodium carboxymethylcellulose, and alginic acid; and a lubricant selected from the